LIANOL: THE NATURAL START!



FERTI IS PART OF THE LIANOL 5 STEP PIG CONCEPT











(3)







(5)

FERTILITY	FARROWING	BIRTH	FARROWING PEN piglet	WEANING
sow	sow	piglet		piglet
Higher farrowing rate	Smooth birth	More energy	Good gut development	Healthy
and more	of vital,	and extra immunity	and digestion	gut
uniform piglets	heavier piglets	right after birth	from day one	around weaning

Lianol 5 step Pig Concept is an innovative feed approach from Ardol.



Ardol BV

Handelsweg 7 6114 BR Susteren The Netherlands T +31 4 64 86 15 05 info@ardol.nl www.lianolconcept.com Available at:





FERTILITY



HIGHER FARROWING

RATE AND MORE

UNIFORM PIGLETS

IN HEAT SOONER; MORE UNIFORM AND MORE PIGLETS



Successful insemination depends on the condition of the sow.

Successful insemination and gestation depend on the condition of the sow, the moment of insemination, and the quality of sperm and egg.

Many sows suffer from an energy deficit at the end of lactation. This causes them to visibly lose weight and it disrupts the metabolism. Good metabolism is important for the development of healthy ova, smooth ovulation and more, viable embryos.

Lianol ferti affects the sow's IGF-1* level, which stimulates the complex fertility process in a natural way. It also boosts metabolism, allowing sows to recover more quickly. This has a positive impact on the hormonal balance of the sow and contributes to the process of ova maturation. They will ovulate sooner, so the sow will come into heat sooner.


LIANOL FERTI; THE BASIS FOR A HIGHER FARROWING RATE

Lianol ferti results in:

- sooner and more visibly in heat
- higher farrowing rate
- more uniform litters
- less adverse effects of heat stress
- less impact of seasonal influences


* IGF-1 is Insulin-like Growth Factor 1, a growth stimulator naturally produced by the body.

Fewer repeated services




Measured variable 235,000 sows

More weaned piglets



Measured variable 235,000 sows

More live-born piglets



Measured variable 235,000 sows

More information: www.lianolconcept.com

LANGE BOOK TO WAR THE STATE OF THE STATE OF

USE OF LIANOL FERTI

Lianol ferti is available as a bolus.

Use:

- Gilts: administer for five days after synchronization treatment
- Second-litter sows: from three days before to three days after weaning
- Older sows: from two days before to two days after weaning